

Synthetic life

(continuation of „The molecular origins of life” SoSe 2023)



NaturalNews.com

WiSe 2024/24

Zbigniew Pianowski

7 lectures (90 min. each) in English

1st lecture: 25th Oct. 2023 (Wed.)

Zoom:

<https://us02web.zoom.us/j/89079835848?pwd=MU83ZXlqa2RWNlFzVXhvNzBocDlGQT09>

Following lecture terms 14:00-15:30, Wed.:

8.11, 15.11., 22.11., 29.11., 6.12. and 13.12.2023 (backup date 20.12.2023).

The most current dates, handouts – on the website:

https://www.ioc.kit.edu/pianowski/99_283.php

and by ILIAS (KIT)

The molecular origins of life

Life is a self-replicating chemical system capable of evolution (NASA, 2009)



Origin of the Universe – stars, planets, elements

Origin of biorelevant monomers – primordial soup

Complex chemical processes on the way to living systems

Protocells and LUCA

Synthetic life



www.genome.gov/about-genomics/policy-issues/Synthetic-Biology

How new functions can emerge from known biological building blocks?

What is Life? What makes it different from just matter?



Everything – living or not – is just chemicals made of atoms.



Every living creature has its code, that makes it grow, reproduce, and change.

DNA turns dust into life.




Fishes swim in water. But what makes fishes alive and not water is the way how the atoms are organized – By the special kind of molecules: DNA – the double helix molecule that houses the genetic alphabet of A, C, G and T, which, in different combinations, can make a flower, or a frog, or you...


20 years ago, Scientists learned to read the creatures' entire DNA sequence, from beginning to the end – the genome



Whole genome sequencing was initially achieved for simple organisms:
bacteria, nematodes, flies and plants...



Haemophilus influenzae
1995



Caenorhabditis elegans
1998



Drosophila melanogaster
2000




Arabidopsis thaliana
2000

... and way up to mammals and human



Mus musculus
2002



Homo sapiens
2004

Human Genome Project (NIH)
Craig Venter – Celera Genomics (private)

With that knowledge, scientists began to tinker...

... to take a glow from a jellyfish...

... and transfer it to a cat...

... or to a rabbit...



To make creatures do what they never did before.

As biologists got better in this, a new kind of science was born – synthetic biology

Definition: *Synthetic Biology*

(also known as *Synbio*, *Synthetic Genomics*, *Constructive Biology* or *Systems Biology*)

„the design and construction of new biological parts, devices and systems that do not exist in the natural world and also the redesign of existing biological systems to perform specific tasks”

Advances in nanoscale technologies – manipulation of matter at the level of atoms and molecules – are contributing to advances in synthetic biology.

What can we do with new tools of synthetic biology?

We can improve what was spelled out for the 3,5 Billion years of evolution.


We can take it beyond reading genomes or editing genomes...

...and start writing genomes.

Our own ideas of what life should be like.

Making creatures drastically different from any that have ever existed.

How could it be done?



Overview of the course

artificial ribozymes and aptamers for efficient catalysis and recognition (SELEX, DNAzymes, foldamers);

unnatural base pairing – expansion of the genetic alphabet;

artificial genetic polymers and oligonucleotide analogues (XNA);


















biosynthetic incorporation of **unnatural aminoacids (UAAs)** into proteins;

enzyme engineering – production of enzymes with unknown or unnatural properties, *ab initio* protein design, directed evolution, theozymes;



Artificial lipid vesicles as models for protocell multiplication;

design of artificial organisms


Artificial genetic polymers




5' -CTTATPPPZZZATAAG-3'
3' -GAATAZZZPPPTATTC-5'




Artificial genetic polymers






Hexitol Nucleic Acid (HNA)




Threose Nucleic Acid (TNA)




Peptide Nucleic Acid (PNA)




Expanded genetic code




Expanded genetic code





a Recoding UGA from stop to Sec



b Reassigning UAG from stop to Pyl



Protein engineering and de novo enzyme design



Introduction

How chemists and biologists are learning from each other?



Greek mythology – introduction to modern molecular biology – chimera, centaur

Introduction

The Central Dogma of the molecular biology – DNA → RNA → proteins


Polymerases and ribosomes - the molecular machines of life

PCR – Polymerase chain reaction – *in vitro* DNA amplification

Recombinant protein production – how to produce a protein in another organism


Protein engineering – how to make desired modifications in proteins

From DNA to proteins




<https://www.youtube.com/watch?v=gG7uCskUOrA>

The Central Dogma: From DNA to proteins




DNA replication



https://www.youtube.com/watch?v=TNKWgcFPHqw&ab_channel=yourgenome


DNA polymerases



Procaryotic DNA Polymerases

Polymerase	Polymerase activity (for all enzymes 5' → 3')	Exonuclease activity
DNA polymerase I	Filling if gap after removal RNA primer, DNA repair, removal of RNA primers	5'→3' and 3'→5'
DNA polymerase II	DNA repair	3'→5'
DNA polymerase III*	Replication, proofreading and editing	3'→5'

*The main enzyme of replication




Nature Reviews | Molecular Cell Biology

Procaryotic DNA Polymerase III

SUBUNITS OF THE DNA POLYMERASE III HOLOENZYME

Subunit	Function	Groupings
α	5'-3' polymerization	Core enzyme: Elongates polynucleotide chain and proofreads
ε	3'-5' exonuclease	
θ	core assembly	
γ	Loads enzyme on template (Serves as clamp loader)	γ complex
δ		
δ'		
χ		
ψ		
β	Sliding clamp structure (processivity factor)	
τ	Dimerizes core complex	

Extremophilic organisms




Brian W. Schaller, Yellowstone Park

Thermus aquaticus is a thermophilic bacteria from hot springs in Yellowstone Park
70°C – optimum, living range: 50-80°C

It is a source of thermostable enzymes

PCR – Polymerase Chain Reaction




Taq polymerase withstands denaturing conditions (hot temperatures) detrimental for most enzymes. *Activity optimum: 75-80°C, half-life at 95°C > 2.5 h*

1990 – Kary Mullis optimized the PCR technique with *Taq* polymerase (1993 Nobel Prize)

<https://www.dnalc.org/view/15475-The-cycles-of-the-polymerase-chain-reaction-PCR-3D-animation.html>

Polymerase chain reaction - PCR



- 1 **Denaturation** at 94-96°C
- 2 **Annealing** at ~68°C
- 3 **Elongation** at ca. 72 °C




Rror




Transcription: DNA → RNA

RNA POLYMERASES IN EUKARYOTES

Form	Product	Location
I	rRNA	Nucleolus
II	mRNA, snRNA	Nucleoplasm
III	5S rRNA, tRNA	Nucleoplasm



Prokaryotic RNA polymerase



Eukaryotic RNA polymerase

Figure 29.1

Biochemistry, Seventh Edition

© 2012 W. H. Freeman and Company


Transcription: DNA → RNA

RNA Polymerase of prokaryotes


Subunit	Function
α, α	Determine the DNA to be transcribed
β	Catalyze polymerization
β'	Bind & open DNA template (unwinding)
ω	Function is not known
σ	Recognize the initiation sites called promoter

<https://www.youtube.com/watch?v=SMtWvDbfHLo>

(a) RNA polymerase binding to promoter




(b) Initiation




Transcription: DNA → RNA

Because transcription start and stop signals are specific to one strand, different genes on the same chromosome can be oriented in different directions. Only one particular strand is ever transcribed for each gene



Transcription: DNA → RNA

How do the start/stop signals look like? For procaryotes (like bacteria *E. coli*):




the transcription start site does not exactly correspond to the codon that is used to initiate translation – base 1 (+1). Transcription begins upstream of this sequence. The RNA between the „base 1” and the start of translation some distance downstream is called the 5' untranslated region or 5' UTR. There is also an untranslated region at the 3' end (3' UTR).

There are two conserved elements in promoter regions of *E. coli*: an element around -10 and an element around -35. These elements can be identified in most *E. coli* promoters.


mRNA processing

In 1977, Phil Sharp (Nobel Prize 1993) hybridized an mRNA to its DNA template and prepared the hybrid molecule for electron microscopy by coating the nucleic acid with a basic protein, then using rotary shadowing to coat the nucleic acid-protein complex.

(a) Micrograph of DNA-RNA hybrid



(b) Interpretation of micrograph




Eucaryotic genes are discontinuous!!!

mRNA processing

The transcript is discontinuous.

There are parts of the DNA template that are not represented in the mRNA.

Eukaryotic mRNA processing




introns are removed (splicing)

When a eukaryotic gene is transcribed, the primary transcript is processed in the nucleus in several ways. The most striking modification is splicing. Parts of the primary transcript, called introns, are spliced out of the mRNA. The remaining segments of mRNA are called exons.

<https://www.youtube.com/watch?v=aVgwr0QpYNE>


mRNA processing

Splicing is facilitated by a ribonucleoprotein complex called the spliceosome. The spliceosome carries out the removal of introns as RNA lariats, joining exons together to make a mature mRNA, as shown below.





Alternative splicing - tropomyosin

Splicing is regulated, with many genes producing multiple isoforms of the same protein that can differ considerably in their amino acid sequence due to alternative splicing. Isoforms of the muscle protein tropomyosin derived from alternative splicing are shown below. While there are some exons common to all isoforms, some isoforms have large protein segments entirely missing from other isoforms



Translation: RNA → proteins




https://www.youtube.com/watch?v=Tfyf_rPWUdY




<https://www.youtube.com/watch?v=kmrUzDYAmEI>

Translation: RNA → proteins

The coupling of transcription and translation in bacteria




Eukaryotic polyribosomes




5' end of the mRNA, because there are shorter protein tails on the ribosomes at that end

Translation: RNA → proteins



Charged serine tRNA

This shows a "charged" serine tRNA, covalently attached to the amino acid serine at its 3' end, with the anticodon paired to a serine codon



Translation: RNA → proteins


Aminoacyl tRNA synthetase

A special set of enzymes "charges" tRNAs, attaching the correct amino acid to particular tRNAs.


A charged tRNA is called an aminoacyl tRNA, so the charging enzymes are more properly called aminoacyl tRNA synthetases.

There is only one aminoacyl tRNA synthetase for each amino acid, even though there can be multiple tRNAs for that amino acid. Each aminoacyl tRNA synthetase is able to recognize all of the tRNAs that need to be charged with the one amino acid that is their specialty.

Amino acids are attached to the hydroxyl (-OH) group at the 3' end of the tRNA through their carboxyl (-COOH) group




Translation: RNA → proteins




Three sites are associated with tRNAs: the A (aminocyl) site, that accepts a new aminoacyl tRNA; the P (polypeptide) site, that holds a tRNA with the growing polypeptide chain; and the E (exit) site that holds an uncharged tRNA ready to exit the ribosome

Translation: RNA → proteins


Translation initiation



1. mRNA binds to small subunit.
Ribosome binding site sequence binds to a complementary sequence in an RNA molecule in the small subunit of the ribosome, with the help of protein initiation factors.




2. Initiator aminoacyl tRNA binds to start codon.



3. Large subunit of ribosome binds, completing ribosome assembly. Translation begins.


Translation: RNA → proteins

Translation elongation




1. Incoming aminoacyl tRNA

New tRNA moves into A site, where its anticodon base pairs with the mRNA codon.



2. Peptide bond formation

The amino acid attached to the tRNA in the P site is transferred to the tRNA in the A site.



3. Translocation

mRNA is ratcheted through the ribosome by elongation factors (not shown). The tRNA attached to the polypeptide chain moves into the P site. The A site is empty.

Translation: RNA → proteins

Translation elongation




4. Incoming aminoacyl tRNA

New tRNA moves into A site, where its anticodon base pairs with the mRNA codon.



5. Peptide bond formation

The polypeptide chain attached to the tRNA in the P site is transferred to the aminoacyl tRNA in the A site.




6. Translocation

mRNA is ratcheted through the ribosome again. The tRNA attached to polypeptide chain moves into P site. Empty tRNA from P site moves to E site, where tRNA is ejected. The A site is empty again.

Translation: RNA → proteins


Translation termination



1. Release factor binds to stop codon.
When translocation exposes a stop codon, a release factor fills the A site. The release factor breaks the bond linking the tRNA in the P site to the polypeptide chain.



2. Polypeptide is released.
The hydrolysis reaction frees the polypeptide, which is released from the ribosome. The empty tRNAs are released either along with the polypeptide or...



3. Ribosome subunits separate.
...when the ribosome separates from the mRNA, and the two ribosomal subunits dissociate. The subunits are ready to attach to the start codon of another message and start translation anew.


Translation: RNA → proteins – the genetic code

nonpolar polar basic acidic (stop codon)


Standard genetic code

1st base	2nd base								3rd base
	U		C		A		G		
U	UUU	(Phe/F) Phenylalanine	UCU	(Ser/S) Serine	UAU	(Tyr/Y) Tyrosine	UGU	(Cys/C) Cysteine	U
	UUC		UCC		UAC		UGC		C
	UUA	(Leu/L) Leucine	UCA		UAA	Stop (Ochre)	UGA	Stop (Opal)	A
	UUG		UCG		UAG	Stop (Amber)	UGG	(Trp/W) Tryptophan	G
C	CUU	(Leu/L) Leucine	CCU	(Pro/P) Proline	CAU	(His/H) Histidine	CGU	(Arg/R) Arginine	U
	CUC		CCC		CAC		CGC		C
	CUA		CCA		CAA	(Gln/Q) Glutamine	CGA		A
	CUG		CCG		CAG		CGG		G
A	AUU	(Ile/I) Isoleucine	ACU	(Thr/T) Threonine	AAU	(Asn/N) Asparagine	AGU	(Ser/S) Serine	U
	AUC		ACC		AAC		AGC		C
	AUA		ACA		AAA	(Lys/K) Lysine	AGA	(Arg/R) Arginine	A
	AUG ^[A]	ACG	AAG		AGG		G		
G	GUU	(Val/V) Valine	GCU	(Ala/A) Alanine	GAU	(Asp/D) Aspartic acid	GGU	(Gly/G) Glycine	U
	GUC		GCC		GAC		GGC		C
	GUA		GCA		GAA	(Glu/E) Glutamic acid	GGA		A
	GUG		GCG		GAG		GGG		G


Expanded genetic code




a Recoding UGA from stop to Sec



b Reassigning UAG from stop to Pyl





Recombinant proteins



Recombinant insulin:

https://www.youtube.com/watch?v=glT8iAqK8NQ&ab_channel=ScienceForStudent


de novo enzyme design




For interested in details:

Introduction to protein design – Part1: https://www.youtube.com/watch?v=0LetJMbU7uY&ab_channel=iBiology



Introduction to protein design – Part2: https://www.youtube.com/watch?v=ZrAwWx7meTk&ab_channel=iBiology




Evolvability of proteins from thermophiles





Aquifex aeolicus (Aa) – thermophilic bacteria



Z. Azuma, T. G. W. Edwardson and D. Hilvert, *Chem. Soc. Rev.*, **2018**, *47*, 3543-3557



B. Wörsdörfer, K. J. Woycechowsky and D. Hilvert, *Science*, **2011**, *331*, 589–592



Overview of the course

artificial ribozymes and aptamers for efficient catalysis and recognition (SELEX, DNAzymes, foldamers);

unnatural base pairing – expansion of the genetic alphabet;

Artificial genetic polymers and oligonucleotide analogues (XNA);

biosynthetic incorporation of **unnatural aminoacids (UAAs)** into proteins;

enzyme engineering – production of enzymes with unknown or unnatural properties, *ab initio* protein design, directed evolution, theozymes;

Artificial lipid vesicles as models for protocell multiplication;

design of artificial organisms